

Timing in the Absence of Clocks: Encoding Time in Neural Network States

Uma R. Karmarkar¹ and Dean V. Buonomano^{2,3,4,*}

¹University of California, Helen Wills Neuroscience Institute, 230 Barker Hall #3190, Berkeley, CA 94720, USA

²Department of Neurobiology

³Department of Psychology

⁴Brain Research Institute

University of California, Los Angeles, Los Angeles, CA 90095, USA

*Correspondence: dbuono@ucla.edu

DOI 10.1016/j.neuron.2007.01.006

SUMMARY

Decisions based on the timing of sensory events are fundamental to sensory processing. However, the mechanisms by which the brain measures time over ranges of milliseconds to seconds remain unclear. The dominant model of temporal processing proposes that an oscillator emits events that are integrated to provide a linear metric of time. We examine an alternate model in which cortical networks are inherently able to tell time as a result of time-dependent changes in network state. Using computer simulations we show that within this framework, there is no linear metric of time, and that a given interval is encoded in the context of preceding events. Human psychophysical studies were used to examine the predictions of the model. Our results provide theoretical and experimental evidence that, for short intervals, there is no linear metric of time, and that time may be encoded in the high-dimensional state of local neural networks.

INTRODUCTION

All forms of sensory processing are ultimately based on decoding the spatial and/or temporal structure of incoming patterns of action potentials. The elucidation of the neural mechanisms underlying the processing of spatial patterns has advanced considerably in the past 40 years. For example, the coding and representation of simple spatial patterns, such as the orientation of a bar of light, are well characterized in primary visual cortex (Hubel and Wiesel, 1962; Ferster and Miller, 2000). Indeed, much has been discovered about the mechanisms underlying the emergence of orientation-selective cells and their role in perception (e.g., Miller et al., 1989; Ferster and Miller, 2000; Gilbert et al., 2000; Schoups et al., 2001; Yang and Maunsell, 2004).

In comparison with spatial stimuli, there is a significant gap in our understanding of how the brain discriminates simple temporal stimuli, such as estimating the duration of time for which a light or tone is presented. Recent studies have begun to examine the neural (Kilgard and Merzenich, 2002; Hahnloser et al., 2002; Leon and Shadlen, 2003) and anatomical (Rao et al., 2001; Lewis and Miall, 2003; Coull et al., 2004) correlates of temporal processing. However, the neural mechanisms that allow neural circuits to tell time and encode temporal information are not clear. Indeed, it has not yet been determined if timing across different time scales and modalities relies on centralized or locally independent timing circuits and mechanisms (Ivry and Spencer, 2004).

Timing is critical in both the discrimination of sensory stimuli (Shannon et al., 1995; Buonomano and Karmarkar, 2002; Ivry and Spencer, 2004; Buhusi and Meck, 2005) and the generation of coordinated motor responses (Mauk and Ruiz, 1992; Ivry, 1996; Meegan et al., 2000; Medina et al., 2005). The nervous system processes temporal information over a wide range, from microseconds to circadian rhythms (Carr, 1993; Mauk and Buonomano, 2004; Buhusi and Meck, 2005). We will focus on the scale of milliseconds and seconds, in which the dominant model of temporal processing is the internal clock model. A prototypical clock model includes an oscillator (pacemaker) that emits pulses that are counted by an accumulator (Creelman, 1962; Treisman, 1963; Church, 1984; Gibbon et al., 1997). Within this framework, the pulse count provides a linear metric of time, and temporal judgments rely on comparing the current pulse count to that of a reference time. This model has proven effective in providing a framework for much of the psychophysical data relating to temporal processing (Church, 1984; Meck, 1996; Rammsayer and Ulrich, 2001). However, electrophysiological and anatomical support for the putative accumulator remains elusive, and mounting evidence indicates that clock models are not entirely consistent with the experimental data (for reviews see Mauk and Buonomano, 2004; Buhusi and Meck, 2005).

A number of alternate models of timing have been suggested (see Discussion; for reviews see Gibbon et al., 1997; Buonomano and Karmarkar, 2002; Buhusi and

Meck, 2005). One such class of models, state-dependent networks (SDNs), propose that neural circuits are inherently capable of temporal processing as a result of the natural complexity of cortical networks coupled with the presence of time-dependent neuronal properties (Buonomano and Merzenich, 1995; Buonomano, 2000; Maass et al., 2002). This framework, based on well-characterized cellular and network properties, has been shown to be able to discriminate simple temporal intervals on the millisecond scale, as well as complex spatial-temporal patterns (Buonomano and Merzenich, 1995; Buonomano, 2000; Maass et al., 2002). Here we examine the mechanisms and nature of the timing in this model and show that it encodes temporally patterned stimuli as single “temporal objects,” as opposed to the sum of the individual component intervals. This generates the counterintuitive prediction that we do not have access to the objective (absolute) time of a given interval if it was immediately preceded by another event. This prediction is tested and confirmed using independent psychophysical tasks. Together, our results provide a mechanistic account of the distinction between millisecond and second timing and suggest that within the millisecond range, timing does not rely on clock-like mechanisms or a linear metric of time.

RESULTS

State-Dependent Networks

An SDN composed of 400 excitatory (Ex) and 100 inhibitory (Inh) recurrently connected integrate-and-fire units was simulated using NEURON. The synapses in the network exhibit short-term forms of synaptic plasticity and both fast and slow IPSPs (see *Experimental Procedures*). Short-term synaptic plasticity (Zucker, 1989) plays a critical role in SDNs by altering the state of the network in a time-dependent fashion after each input, which in turn produces time-dependent neuronal responses. In essence, in the same manner that long-term plasticity may provide a memory of a learning experience (Martin et al., 2000), SDNs use short-term synaptic plasticity to provide a memory trace of the recent stimulus history of a network (Buonomano, 2000).

The functional properties of an SDN can be understood if we consider the sequential presentation of two brief and identical events (e.g., two auditory tones) 100 ms apart (Figure 1A). When the first event arrives in the network, it will trigger a complex series of synaptic processes resulting in the activation of a subset of neurons. When the same event is repeated 100 ms later, the state of the network will have changed from S_0 to S_{100} . Due to the time-dependent changes in network state (imposed by short-term synaptic plasticity), the population response to the second stimulus inherently encodes the fact that an event occurred 100 ms before. In this fashion the network implements a temporal-to-spatial transformation—i.e., the presence, absence, or number of spikes from a given subset of neurons will depend on the temporal structure of the stimulus. The model

is stochastic in the sense that determining which neurons will be interval sensitive is a complex function of the network's random connectivity, assigned synaptic strengths, and short-term plasticity (Buonomano, 2000). Once time is encoded in a spatial code, it can be read out by a set of output neurons (see below; Buonomano and Merzenich, 1995; Buonomano, 2000; Maass et al., 2002; Knüsel et al., 2004).

In this model, there is no explicit or linear measure of time like the tics of an oscillator or a continuously ramping firing rate (see *Discussion*; Durstewitz, 2003). Instead, time is implicitly encoded in the state of the network—defined not only by which neurons are spiking, but also by the properties that influence cell firing, such as the membrane potential of each neuron and synaptic strengths at each point in time. Thus, even in the absence of ongoing activity, the recent stimulus history remains encoded in the network. The simulation in Figure 1 consists of 500 neurons and a total of 12,200 synapses, allowing us to define the network's state in 12,700-dimensional space. Since the state of the network ultimately determines the response to the next input, we can think of its evolving trajectory through this space as encoding time. Principal component analysis was performed to provide a visual representation of this trajectory (see *Experimental Procedures*). In response to a single stimulus, the first three principal components establish a rapidly evolving neural trajectory through state-space, followed by a much slower path settling back toward the initial state (Figure 1B). When a second event is presented at $t = 100$ ms, it produces a perturbation in state-space different from the $t = 0$ event (Figure 1C). Similarly, additional presentations of the same stimulus at varying delays would continue to produce cumulative changes in network state.

The time it takes for the network to return to its initial state—its reset time—is a function of the longest time constants of the time-dependent properties. For short-term synaptic plasticity, this is on the order of a few hundred milliseconds (Zucker, 1989; Markram et al., 1998; Reyes and Sakmann, 1998). The dynamics of short-term plasticity must run its course; thus, the network cannot return to its initial state on command. As addressed below, this property has important implications for temporal processing.

Temporal Objects

An important feature of SDNs is that they naturally extend beyond simple interval discrimination to the processing of complex temporal sequences. This is due to the cumulative nature of changes in network state (Buonomano and Merzenich, 1995; Maass et al., 2002). However, potential weaknesses in SDNs arise because of both the absence of an explicit metric of time and their sensitivity to changes in initial state.

To examine these issues we investigated the ability of the network to discriminate between 100 and 200 ms intervals (we will use the notation [100] × [200] ms), as well as two simple patterns that contain these intervals, namely




Figure 1. State-Dependent Network Simulation

(A) Voltage plot of a subset of neurons in the network. Each line represents the voltage of a single neuron in response to two identical events separated by 100 ms. The first 100 lines represent 100 Ex units (out of 400), and the remaining lines represent 25 Inh units (out of 100). Each input produces a depolarization across all neurons in the network, followed by inhibition. While most units exhibit subthreshold activity, some spike (white pixels) to both inputs, or exclusively to the first or second. The Ex units are sorted according to their probability of firing to the first (top) or second (bottom) pulse. This selectivity to the first or second event arises because of the difference in network state at $t = 0$ and $t = 100$ ms.

(B) Trajectory of the three principal components of the network in response to a single pulse. There is an abrupt and rapidly evolving response beginning at $t = 0$, followed by a slower trajectory. The fast response is due to the depolarization of a large number of units, while the slower change reflects the short-term synaptic dynamics and slow IPSPs. The speed of the trajectory in state-space can be visualized by the rate of change of the color code and by the distance between the 25 ms marker spheres. Because synaptic properties cannot be rapidly “reset,” the network cannot return to its initial state (arrow) before the arrival of a second event.

(C) Trajectory in response to a 100 ms interval. Note that the same stimulus produces a different fast response to the second event. To allow a direct comparison, the principal components from (B) were used to transform the state data in (C).

a 100 or 200 ms interval preceded by a 150 ms interval ([150; 100] and [150; 200]). We calculated the information each neuron in the network contains for the discrimination of both sets of stimuli. Mutual information was determined based on the number of spikes in each neuron (see *Experimental Procedures*). The neurons containing information for the [100] \times [200] and the [150; 100] \times [150; 200] discriminations fall in largely nonoverlapping populations (Figure 2A). This occurs even though the discrimination could in principle be based on the same [100] \times [200] interval. Since the individual intervals are encoded in the context of the whole stimulus, the network cannot recognize that the [100] and [150; 100] patterns share a common feature. Nevertheless, it can discriminate between all four stimuli (Figure 2B). Each stimulus is coded as a distinct temporal object regardless of its component features.

Reset Task

The prediction that emerges from the model is that if a distractor precedes a 100 ms target interval at random intervals, discrimination of the target should be impaired in comparison to a 100 ms interval with no distractor (or

one preceded by a fixed distractor). This prediction was examined using psychophysical studies. We designed a task (Figure 3A) in which each trial consisted of a randomly interleaved presentation of a single two-tone (2T) or three-tone (3T) stimulus, and participants were asked to judge the interval between the last two tones. In the 3T case the first tone acts as a distractor. By independently and adaptively varying the intervals, discrimination thresholds were calculated for the 2T and 3T tracks (see *Experimental Procedures*). The randomly interleaved—and thus unpredictable—presentation of the 3T stimuli also ensured that the subjects did not adopt strategies to ignore the distractor. The standard interval (SI) was presented at the beginning of a trial and maintained implicitly as a result of feedback to each response (Grondin and Rammsayer, 2003; Karmarkar and Buonomano, 2003). Subjects were asked to judge whether the target interval was shorter or longer than the standard. Two classes of distractors, fixed (FIX) and variable (VAR), were examined. In the FIX condition, the distractor was always presented at a fixed interval before the target interval. In the VAR condition, the distractor was presented at a range of times ($\pm 50\%$ of the standard).




Figure 2. Encoding of Temporal Patterns

(A) Information per neuron. The blue trace displays the mutual information that each Ex unit provides for the discrimination of a 100 versus 200 ms interval (sorted). The red line shows the information for the same intervals preceded by a 150 ms interval; that is, discrimination of the pattern [150; 100] versus [150; 200]. While individual neurons contain significant information for both stimuli, a different population of neurons encodes each one.

(B) Discrimination of all four stimuli. All Ex units were connected to four output neurons trained to recognize the network activity produced by the last pulse of all four stimuli. Average responses were calculated from six independent (different random number generator seeds) simulations. Note that a mutual information measure based on total spike count to each stimulus, as in (A), would introduce a confound because the number of spikes is also a function of the number of events (see [Experimental Procedures](#)). Each group of four bars represents the responses of the four output neurons.

This task was termed the “Reset task” based on the unique constraints it places on the temporal encoding mechanisms. If a subject were using a simple stopwatch strategy, he or she would have to start the stopwatch at the first tone, even though it is irrelevant in the 3T trials. The true role of the second tone can only be determined retroactively by the presence or absence of a third tone. With a stopwatch, one approach could be to quickly record the time at t_2 and then reset the watch. Alternately, the time at t_2 and t_3 could be noted and then t_2 subtracted from t_3 to obtain the interval between the second and third tones. We will refer to the first strategy as a clock reset mechanism and the second as temporal arithmetic. Both can be implemented with internal clock models, either because the accumulator could be reset, or because the presence of a linear temporal metric would allow for temporal arithmetic. Both clock-based models predict that performance on the 2T and 3T tracks should be similar in both the FIX and VAR conditions because the predictability of the distractor should not affect the encoding of t_1 - t_2 and t_2 - t_3 .

In contrast, in the SDN model, a reset strategy cannot be implemented because short-term plasticity cannot be reset on cue. Temporal arithmetic cannot be performed due to the absence of a linear metric of time. SDNs predict that performance on the FIX condition will be similar for the 2T and 3T stimuli because the feedback at the end of each trial can be used to establish consistent states on which to build internal temporal representations for both stimuli. However, they also predict an impaired performance in the 3T-VAR trials compared with the 2T or FIX conditions since the state of the network will not be reproducible across trials.

Subjects were first tested with a target interval of 100 ms (SHORT). Consistent with previous studies, thresholds for the 2T conditions were in the range of 20% of the target ([Wright et al., 1997](#); [Karmarkar and Buonomano, 2003](#)). A two-way analysis of variance (ANOVA) revealed a significant interaction between conditions (FIX \times VAR) and tone number (2T \times 3T; $F = 57.75$; $n = 15$; $p < 0.0001$), demonstrating a dramatic impairment in the 3T-VAR condition only ([Figure 3B](#)). Indeed, the threshold in the 3T-VAR condition for a 100 ms interval was similar to that observed in independent (2T only) experiments on a 200 ms interval (46 ± 3.4 ms versus 45 ± 7 ms; data not shown). Thus, under the SHORT condition, the psychophysics supported the predictions of the SDN. In contrast, when the Reset task involved a target of 1000 ms (LONG), there was no effect of the variable distractor, as evident in the lack of interaction in the ANOVA ([Figure 3C](#); $F = 0.087$; $n = 12$; $p > 0.5$). Importantly, the point of subjective equality (PSE) was approximately equal to the target intervals in both the SHORT and LONG experiments, independent of the presence or absence of the distractor in both the FIX and VAR conditions ([Figures 3D and 3E](#)). Therefore, a memory component of the task cannot account for the differences observed between the two target lengths.

The specific effect of the variable distractor on the SHORT group is consistent with the prediction of the SDN model. It is unlikely that this result is due to effects such as the increased uncertainty caused by the variable distractor, as the same degree of uncertainty was present in the LONG trials without an accompanying timing impairment. Additionally, the randomly interspersed presentation of the 2T and 3T stimuli ensures the same level of uncertainty for both stimuli (in both conditions), but the 2T-VAR performance was not affected. However, to further examine the general psychophysical effects of a variable distractor, we conducted two additional controls. The first was a task in which the distractor interval was 100 (FIX) or 50–150 ms (VAR) coupled with a 1000 ms target (Short-Long). In addition, subjects performed a frequency discrimination task in which the target frequency was preceded by a tone either at a fixed or variable interval (see [Experimental Procedures](#)). Neither the Short-Long [$F = 0.18$; $n = 10$; $p > 0.5$] or frequency [$F = 0.23$; $n = 14$; $p > 0.5$] experiments revealed a decrement in performance produced by the variability of the distractor ([Figures 4A and 4B](#)).




Figure 3. Reset Task: a Variable Distractor Impairs Discrimination of a Short, But Not a Long, Interval

(A) Reset task. Top rows represent the standard 2T interval discrimination task in a single stimulus protocol. Subjects are asked to press different mouse buttons if they judged the interval to be short (S) or long (L). The feedback across trials results in the creation of an internal representation of the target interval. Bottom rows represent the 3T task in which a distractor is presented at a fixed or variable (dashed) interval across trials.

(B) Thresholds for the 100 ms (SHORT) Reset task. (Left) Thresholds for the 100 ms 2T interval discrimination (open bars) and the 100 ms interval preceded by a distractor presented at the same interval across trials (3T-FIX; gray). (Right) Threshold for the standard 100 ms task (open) and 3T task in which the distractor was presented at variable intervals across trials (3T-VAR; gray). Error bars = SEM. The asterisk represents a significant difference from the other three groups.

(C) Reset task (represented as in A), using a 1000 ms (LONG) target interval. Neither of the main effects nor the interaction was significant.

(D and E) Point of subjective equality (PSE) values for the same experiments shown in (B) and (C), respectively. The PSE was not significantly different from the target intervals of 100 (D) and 1000 ms (E) in any condition.

Effect of the Interstimulus Interval on Performance

It is important to rule out the possibility that the impairments observed in the Reset task were not produced by some complex interaction between uncertainty and the intervals being judged, or that the distractor in the FIX condition was serving as a reference interval (see Discussion). Thus, we examined the prediction of the SDN model using a second independent psychophysical test. The SDN model predicts impaired performance under conditions when the network state at the time of the target stimulus varies across trials. This condition can also be produced by insufficient reset time before the next stimulus is presented. To test this directly, we examined performance on a traditional two-interval two-alternative forced-choice task (Wright et al., 1997) in which the interstimulus interval (ISI) was varied. In this paradigm, subjects heard both the 100 ms target and a longer comparison interval, then made a judgment as to whether the longer stimulus occurred first or second. We presented the two intervals with a mean ISI of either 250 or 750 ms. Since experimental data suggests that short-term plasticity operates on the time scale of a few hundred milliseconds (Markram et al., 1998; Reyes and Sakmann, 1998), the state-dependent

model predicts that the network will not have completely returned to its initial state in the ISI_{250} condition, thus impairing temporal discrimination. Indeed, a comparison of the ISI_{250} to the ISI_{750} condition showed a significant decrease in performance for the shorter ISI [$t = 3.53$; $n = 10$; $p < 0.01$] (Figure 5A). Subjects also performed a frequency discrimination task under the short and long ISI conditions, for which they reported if the tone pitch was higher for the first or second stimulus. There was no difference between the two conditions [$t = 0.53$; $n = 10$; $p > 0.5$] (Figure 5A), indicating that the effect of the shorter ISI was specific to time discrimination.

The state-dependent framework predicts that the two intervals are more difficult to compare, resulting in higher temporal discrimination thresholds, because their state-space trajectories have different starting points which vary from trial to trial. The total length of time from the first tone of the first stimulus to the first tone of the second is determined by the exact duration of the ISI ($250\% \pm \leq 25\%$). As a result, the variability in the initial state for the second stimulus is caused by the first—the first interval interferes with the second. However, if the target and comparison stimuli were presented at the same ISI, but




Figure 4. Control Interval and Frequency Discrimination Tasks

(A) Short-Long Reset task. The variable distractor in these trials was between 50–150 ms, and the target interval was 1 s. When a short unpredictable distractor preceded a long target interval, there was no effect of whether the distractor was fixed or variable.

(B) Frequency task. A tone was presented in the absence of a distractor (open bars) or in the presence of a distractor tone presented at a fixed (gray bar, left) or variable (gray bar, right) interval before the target tone. Conventions as in Figure 3.

to different local networks, the impairment produced by the short ISI should be decreased or absent. To examine this prediction, we took advantage of the known tonotopic organization of the auditory system. We performed interval discrimination tasks under two experimental conditions: (1) as above, a 100 ms standard and a comparison (100 + Δ T ms) played at 1 kHz at ISI₂₅₀ and ISI₇₅₀; (2) a similar condition except that one of the stimulus intervals was played at 4 kHz and the other at 1 kHz. Replicating the




Figure 5. Short Interstimulus Intervals Impair Interval, but Not Frequency, Discrimination

(A) Bars on the left show the thresholds for a two-interval two-alternative forced-choice discrimination with a 100 ms target. When the interval between the stimuli was short (250 ms), performance was significantly worse compared with that in the long ISI condition (750 ms). In contrast, performance on a frequency discrimination task was unaltered by the ISI.

(B) Bars on the left illustrate the results for short (250 ms) and long (750 ms) ISI when both the standard and comparison intervals were presented at the same frequency. Bars on the right represent the interval discrimination thresholds when the standard and comparison stimuli were presented at different frequencies. We believe the difference in absolute interval discrimination between both studies (right bars in A and B) reflects interference between the different task and stimulus sets in both studies, as well as the inherent subject variability observed in timing tasks.

above results, Figure 5B shows that there was a significant increase in the threshold of the ISI₂₅₀ compared with the ISI₇₅₀ tasks [$t = 6.85$; $n = 9$; $p < 0.001$] in the same frequency condition. However, using different frequencies for the standard and comparison intervals eliminated any impairment in performance on the short ISI [$t = 0.85$; $n = 9$; $p > 0.3$].

Interval Discrimination despite Differences in Initial State

While the insufficient reset time in the above experiments (Figure 5A) impaired discrimination thresholds, it did not entirely prevent subjects from performing the task. We were thus interested in returning to the theoretical model to determine how performance varied as a function of ISI and whether some degree of timing was still possible with only a partial reset of the network. First, the trajectory of the network in state-space was calculated in response to two 100 ms intervals separated by a 250 or 750 ms ISI. As shown in (Figure 6A), a 750 ms ISI allows the network to return to a point very close to its “naive” initial state. As a result, the trajectory produced by the second stimulus closely traces that produced by the first one. In contrast, for the 250 ms ISI, the network does not return to the neighborhood of the initial state, and its trajectory for the second interval is significantly different. Measures of these distances are presented in Figure 6B.

To quantify the effect of initial state on interval discrimination, output units were trained to discriminate 100 ms from other intervals in the range of 50–150 ms. We then determined the ability of the model to perform this discrimination when the comparison intervals followed the 100 ms target by ISIs that varied from 250–750 ms. Performance worsened with decreasing ISIs (Figure 6C). Importantly, performance changed in a graded manner, indicating that the reset effect is not expected to be all or none. Thus, the behavior of the theoretical model is consistent with the results seen in the human psychophysical data.

DISCUSSION

The standard model of temporal processing postulates a single centralized internal clock, which relies on an oscillator and an accumulator (counter) (Creelman, 1962; Treisman, 1963; Church, 1984; Grondin, 2001). The clock concept is generally taken to imply that the passage of time is counted in units that can be combined or compared linearly. In contrast, SDN models propose that for spans on the scale of tens to hundreds of milliseconds, time may be represented as specific states of a neural network. Within this framework, a 50 ms interval followed by a 100 ms interval is not encoded as the combination of the two. Instead, the earlier stimulus interacts with the processing of the 100 ms interval, resulting in the encoding of a distinct temporal object. Thus, temporal information is encoded in the context of the entire pattern, not as conjunctions of the component intervals.




Figure 6. Dependence of the State-Dependent Network on Initial State

(A) Trajectory of the same network shown in Figure 1 and Figure 2, in response to two 100 ms intervals separated by a 250 (A1) or 750 (A2) ms ISI. Note that the trajectories under the 750 ms ISI are much closer to overlapping than they are in the 250 ms condition. Arrows indicate the times of the onset of the second interval.

(B) Distance matrix. The diagonal represents the distance in Euclidean space between the trajectories shown in (A1) and (A2) starting at 0. The distance is zero until the onset of the second tone (the noise “seed” was the same for both simulations). The secondary diagonals permit the visualization of the distances between two trajectories shifted in time. This allows the comparison of the trajectory starting at the onset of the second interval (for the 250 ms ISI) with that of the first interval (blue rectangle and blue line in lower panel), or the second interval of the 750 ms ISI with the first interval (red rectangle and red line in lower panel). These distances, shown in the lower panel, allow for quantification of the effect of the network not returning to its initial (resting) state before presenting the next stimulus. Note that while the initial distance is lower in the 750 ms ISI, it is not zero.

(C) Percent correct performance of networks trained to discriminate two intervals separated by varying ISIs. Average data from four stimulations. Output units were trained to discriminate intervals ranging from 50–150 ms. Performance was then tested by examining generalization to these same intervals when presented at varying ISIs after the presentation of a 100 ms interval. Results for the 100 × 150 ms discrimination are shown. Performance is highly dependent on the initial state of the network.

State-Dependent Networks and the Reset Task

SDN models propose that timing is a ubiquitous component of neural computations, and that local cortical circuits are inherently capable of processing both temporal and spatial information (Buonomano and Merzenich, 1995; Buonomano, 2000; Maass et al., 2002). In these models timing relies on mechanisms analogous to using the evolving state of a physical system—like the ripples on the surface of a lake—to tell time. However, as shown here (Figure 1 and Figure 2), reliance on the state of a complex system to tell time creates potentially serious limitations due to the resulting dependence on the initial state and the lack of a linear metric of time.

Interestingly, our psychophysical results reveal the same limitations—interval discrimination is impaired by the presence of a distractor that appears at unpredictable times. However, interval discrimination was not altered if the distractor occurred at a fixed time prior to the target. Thus, internal representations of the target interval can develop across trials for the 2T and 3T-FIX stimuli, but not for

the target interval of the 3T-VAR stimuli. This is because the state of the system at the onset of the second tone is variable. The impairment in the 3T-VAR condition is not due to the unpredictability of the distractor’s presence itself; since the 2T and 3T stimuli are randomly intermixed, the unpredictability is the same under all conditions. Rather, the impairment in the 3T-VAR condition is limited to the predictability (consistency) of the *interval* of the distractor.

An alternate interpretation of the 3T-VAR impairment is that in the 3T-FIX condition, the distractor interval served as a reference cue for the target interval. The two-interval discrimination task, in which both a standard and comparison interval are presented on each trial, was used to rule out this possibility (Grondin and Rousseau, 1991; Rammeyer, 1999; Wright et al., 1997). Performance was impaired if the time between the stimuli was 250 ms, but not 750 ms (Figure 5A). It could be argued that the impairment for short ISIs reflects a difficulty in segmenting or attending to rapidly presented stimuli. We find this

interpretation unlikely since performance on the short and long ISI conditions did not differ when the two intervals were presented at different frequencies.

The influence of preceding stimuli on temporal judgments is surprising because much of the timing performed by the nervous system on the scale of hundreds of milliseconds is based on a continuous barrage of incoming stimuli, such as speech or Morse code recognition. The subjects in the current study were naive; thus, a critical issue relates to the effect of learning. We speculate that training would allow subjects to improve their discrimination of intervals independent of temporal context. Indeed, SDN models do not predict that spatial-temporal patterns preceded by other events are impossible to process. Rather, they propose that there must be previous exposure to a large number of instances of the stimuli so that a correspondence between the target information in a number of different contexts can be learned.

Clock Models

The standard clock models predict a linear metric of time, which implies that the clock can time the sequential intervals independent of the presence of a variable distractor across trials. However, most of these models do not explicitly address the issue of the clock reset properties. Thus, it seems reasonable to consider whether a clock with some state-dependent properties could account for the impaired timing of short ISIs or intervals with a distractor. For example, one could assume that resetting or reading the time of the clock is state-dependent, and thus, the reset process could inject noise into the system or be delayed dependent on the initial state.

There are two aspects of our results which could argue against a state-dependent clock mechanism. First, though a state-dependent reset of a centralized clock could explain impaired timing in the short ISI condition (Figure 5A), it would not predict the lack of impairment in the short ISI condition with different frequencies (Figure 5B). The second issue concerns the specificity of the reset problem. In our Reset experiments (Figure 3), a clock would be started by the first tone and stopped and reset (restarted) by the second. The third tone would again stop the clock. As mentioned above, a state-dependent reset would take time or inject noise into the process, and impair the 3T-VAR sequence compared with the 2T one. However, such a clock would also be expected to impair timing of 3T stimuli in the FIX condition. In both cases, the second tone would stop and reset the clock, because there is a 50% chance that the second tone would be the end versus the beginning of the target interval. This prediction is counter to our psychophysical results. One might then propose the use of multiple clocks, in which the first tone activates a primary clock, the second tone activates a secondary clock (and stops the first), and the third tone stops the second clock. This explanation would correctly suggest that timing is not impaired in the FIX condition, but would also hold for the VAR, again violating the dissociation found in our data.

Nevertheless, we cannot eliminate the possibility that there exists a set of assumptions which can enable clock models to account for the observed millisecond timing results. However, we argue that the SDN model provides the most parsimonious explanation of the current psychophysical data on the processing of short intervals.

Other Models of Temporal Processing

A number of other mechanistic models have been put forth to account for measuring and encoding time. These include climbing firing rate models (Durstewitz, 2003; Reutimann et al., 2004), multiple oscillator models (Miall, 1989; Matell and Meck, 2004), and those based on ongoing network dynamics (Medina and Mauk, 2000; Buonomano, 2005). The latter focus primarily on generating appropriately timed motor responses and will not be discussed here.

The climbing or ramping firing rate models suggest that, like many other stimulus features, time is encoded in the firing rate of neurons. Experimentally it is established that some cortical neurons undergo a more or less linear ramping in their firing rate over time (Niki and Watanabe, 1979; Brody et al., 2003; Leon and Shadlen, 2003). In their simplest form climbing models propose that firing rate represents a linear metric of absolute time. However, recent data suggests that, at least in some cases, these neurons are coding expectation rather than absolute time (Janssen and Shadlen, 2005). Climbing rate models have been discussed primarily in relation to timing of intervals or durations; how they would account for timing of temporal patterns has not yet been addressed. Thus, their predictions for our tasks are not immediately clear. For the Reset task it could be argued that ramping would begin at the first stimulus. Time could be read out in the firing rate at the onset of the second and third tone, assuming activity is not reset by the second tone. However, climbing models would not predict the dramatic impairment observed in the 3T-VAR condition or the effect of short ISIs. We would speculate that ramping firing rates are likely to play an important role in the timing of expected motor responses, but less likely to be involved in the timing of rapid sensory stimuli, particularly for complex tasks such as speech or interpretation of Morse code.

The multiple oscillator model suggests that time is encoded in a population of oscillators with different base frequencies (Miall, 1989; Matell and Meck, 2004; Buhusi and Meck, 2005). Time can be read out by a set of coincidence detectors. This model has the advantages of not requiring an accumulator and being capable of timing multiple consecutive intervals once the oscillators have been triggered. However, how this model will behave in the tasks examined here is again dependent on its assumptions. If each event does not reset the oscillators, this model would be expected to produce a decrease in performance in the 3T-VAR condition, consistent with our results. However, it would not necessarily predict the decrease in performance observed with the short ISIs observed in Figure 5, since its reset mechanisms could be all or none. Furthermore, this model posits that timing is centralized. Thus,

it would not predict that any effect of a short ISI would be dependent on whether the frequencies of the comparison stimuli were the same. We would concur that a multiple oscillator model could contribute to timing in the range of seconds (Matell and Meck, 2004; Buhusi and Meck, 2005), but would argue that it is unlikely to account for the timing on the scale of a few hundred milliseconds.

Millisecond versus Second Timing

Timing in both the range of milliseconds and seconds has often been considered to rely on the same underlying mechanisms (Church, 1984; Macar et al., 2002). The results described here demonstrate qualitative differences in the processing of short and long intervals. Unlike the millisecond range, timing of intervals lasting one second or longer appears consistent with mechanisms that generate a linear metric of time. For a 1 s target subjects could accurately judge the first or second of two consecutive intervals in the Reset task, even though they did not know a priori which was the target. Performance was also independent of both fixed and variable distractors preceding the target interval. This implies that subjects could independently keep track of the objective time of two sequential second-long intervals and implies the presence of a linear metric of time. As described above, two simple strategies that a standard clock model could utilize to perform this task are resetting a clock at the second tone, or contributing values to the performance of temporal arithmetic. For the long intervals we did not observe any decrease in timing accuracy in the 3T versus 2T stimuli. We would suggest that this observation is more consistent with the temporal arithmetic scenario. Specifically, that timing on the order of seconds relies on a linear metric of time, and that the second of two consecutive intervals can be calculated by subtracting the first interval from the final count.

The theoretical framework and psychophysical results described here, together with previous psychophysical (Rammsayer and Lima, 1991), pharmacological (Rammsayer, 1999), and imaging studies (Lewis and Miall, 2003), support the existence of distinct loci for subsecond and second processing. The precise boundary between these forms of temporal processing cannot yet be established. However, it seems likely that they are highly overlapping, and that timing in intermediary ranges (e.g., 400–800 ms) may be accurately performed by both the mechanisms underlying time perception and time estimation. Based on the time constants of short-term synaptic plasticity and other time-dependent neural properties, we suggest that the SDN model is limited to intervals below 500 ms. Additionally, even within a specific time scale, there may be multiple mechanisms contributing to timing, and thus the above models are not mutually exclusive.

Relation to Previous Psychophysical Data

A comprehensive model of temporal processing should provide a detailed description of the neural mechanisms

underlying timing, generate novel testable predictions, and account for existing experimental data. Two of the most robust features of temporal processing determined experimentally relate to the scalar property and the role of attention in subjective time estimation. The scalar property refers to the observation that the ratio of the absolute criterion interval and the standard deviation of temporal estimates tends to be constant for long intervals (Gibbon, 1977; Gibbon et al., 1997; Buhusi and Meck, 2005). However, this is not the case for interval discrimination in the range of a few hundred milliseconds (Wright et al., 1997; Mauk and Buonomano, 2004). Thus, we examined how performance scales with short intervals in the SDN model. Results showed that, consistent with the human psychophysical data, temporal resolution is proportionally worse for short intervals (see Figure S1 in the Supplemental Data).

Attention has been widely reported to alter estimates of time in the range of seconds (Hicks et al., 1976; Macar et al., 1994; Brown, 1997; Coull et al., 2004). Internal clock models can account for attention-dependent effects in the second range by assuming a gating mechanism that controls the number of events generated by the oscillator that are counted by the accumulator (Meck, 1984; Zakay and Tsal, 1989). In contrast, on the shorter time scale, divided attention or cognitive load does not appear to specifically alter temporal judgments (Rammsayer and Lima, 1991; Lewis and Miall, 2003). Therefore, the SDN model would be expected to be fairly insensitive to shifts in attention. However, recent studies have revealed that temporal distortions of short intervals can be produced by saccades or stimulus features (Morrone et al., 2005; Johnston et al., 2006). These studies suggest that on short scales, timing is local, and are generally consistent with the SDN model that predicts that temporal processing could occur in a number of different cortical areas on an as-needed basis.

Conclusion

We propose here that cortical networks can tell time as a result of time-dependent changes in synaptic and cellular properties, which influence the population response to sensory events in a history-dependent manner. This framework is applicable to the processing of simple intervals as well as more complex spatial-temporal patterns, and does not invoke any novel hypothetical mechanisms at the neural and synaptic level. Additionally, we propose that timing is not centralized, and can potentially occur locally at both early and late stages of cortical processing. The psychophysical experiments examined here emerged as a direct prediction of this model, and the results are supportive of this general framework. However, establishing the neural basis for timing will ultimately require the accumulation of converging evidence from a number of different fields; of particular relevance will be the use of more complex temporal stimuli in conjunction with in vivo electrophysiology to determine if the population response to ongoing sensory events also contains information about the preceding stimuli.

EXPERIMENTAL PROCEDURES

Neural Network Simulations

The simulated network was composed of 400 Ex and 100 Inh recurrently connected Hodgkin-Huxley units (Buonomano, 2000). Excitatory neurons were randomly interconnected with a probability of 0.2. The mean synaptic weights were adjusted so that neurons responded with zero to three spikes to a short stimulus, as is typical for primary sensory cortex in awake animals (Brody et al., 2002; Wang et al., 2005). Short-term dynamics of excitatory synapses were simulated according to Markram et al. (1998). Short-term synaptic plasticity of Ex → Ex synapses was facilitatory, based on experiments suggesting that paired-pulse facilitation is present in adult cortex (Reyes and Sakmann, 1998; Zhang, 2004). The mean U (utilization), τ_{rec} (recovery from depression), and τ_{fac} (facilitation) parameters were 0.25, 1 ms, and 100 ms, respectively. All three values were randomly assigned using a normal distribution with an SD of 20% of the mean. Short-term plasticity IPSPs in the form of paired-pulse depression was implemented as previously described (Buonomano, 2000).

Mutual Information and Network Readout

Mutual information was calculated using the total number of spikes in response to a stimulus, thus providing an assumption-independent estimate of the amount of information available (Buonomano, 2005). For the discrimination between stimuli with different numbers of pulses (Figure 2A), training of the output units was based on previously described supervised learning rules (Buonomano, 2000; Maass et al., 2002) using only the pattern produced by the last pulse. Training was performed on a set of 25 stimulus presentations and tested on 10 novel test presentations. In the stimulations shown in Figure 6, the outputs were trained to discriminate pairs of intervals (100 ms versus intervals ranging from 50–150 ms). In each case the shortest interval was defined as the short stimulus and the longest as the long stimulus.

Principal Component Analysis

The data set was comprised of the voltage of all Ex and Inh neurons, as well as the synaptic weights (which were time-varying) of excitatory and inhibitory synapses. To reduce the dimensionality of the data set, only 20% of all synaptic weights were used. The data were normalized and the principal components were calculated using the PRINCOMP function in Matlab. Although the dimensionality is very high, the dimensions are highly correlated during the silent period between events (if one cell is hyperpolarized, most cells are hyperpolarized). As a result, the first three principal components can account for a significant amount of the total variability (approximately 75% in Figure 1B). As expected, these components do not account well for the actual response to each event, which is dominated by highly nonlinear dynamics.

Psychophysics

Subjects consisted of graduate and undergraduate students between the ages of 18 and 30 from the UCLA community. All subjects had normal hearing.

Reset Task

These experiments were based on a single-stimulus two-alternative forced-choice protocol as described previously (Karmarkar and Buonomano, 2003). A within-subject design was used; thus, each subject performed the two distractor conditions (FIX and VAR) with each condition having two tracks (2T and 3T). Sessions of the FIX and VAR conditions were given on alternating days over a 1 day period (counterbalanced). Each block within a session consisted of 120 trials: 60 2T and 60 3T. Each tone (1 kHz) was 15 ms in duration and included a 5 ms linear ascending and descending ramp. In the FIX condition, a distractor tone was presented at a fixed interval equal to that of the SI prior to the target. In the VAR condition the distractor occurred before the target at an interval uniformly distributed between

$SI \pm (0.5 \times SI)$. Thresholds for the 2T and 3T tracks were obtained by presenting the target interval as $SI \pm \Delta T$, where ΔT varied adaptively according to a three-down one-up procedure (Levitt, 1971; Karmarkar and Buonomano, 2003). Threshold was defined as two times the mean of the reversal values, which corresponds to a 79% correct performance level.

In each trial subjects made a forced choice decision as to whether the stimulus seemed shorter or longer than the target interval by pressing one of two buttons on a computer mouse. They were provided with immediate visual feedback. All stimuli were generated in Matlab and presented through headphones.

The 2T and 3T stimuli were randomly interleaved to ensure that subjects did not develop a strategy that involved ignoring the distractor tone. Additionally, the simultaneous measure of performance on a conventional 2T task and a task with the presence of a distractor provided a control for nonspecific effects such as difficulty of the overall task, attention, and memory. Target intervals were either 100 or 1000 ms.

A similar protocol to the one used above was also used for the frequency discrimination task. Rather than adaptively varying the interval of the tones, their frequency was varied according to $F \pm \Delta F$ (where F, the target frequency, was 1 kHz). Tone duration was 25 ms.

Two-Interval Forced Choice Procedure

In this task subjects were presented with two intervals on each trial: an SI and the comparison interval (standard + ΔT) (Allan, 1979; Karmarkar and Buonomano, 2003). Subjects were asked to press one of two buttons depending on whether they judged the first stimulus or the second interval to be longer. The SI was 100 ms, and the ISIs for the short and long ISI conditions were 250 and 750 ms, respectively.

The frequency task in the ISI experiments used the same type of stimuli, but shifted the frequency of both tones of the comparison stimulus. Note that in contrast to the single stimulus protocol, subjects could reference the target frequency on each trial as opposed to developing an internal representation of it across trials. We believe this difference, together with the absence of a distractor, is responsible for the improvement in the frequency thresholds as compared with the Reset task. All subjects performed all four tasks in a counterbalanced manner.

Statistics

In the Reset task, the key analysis was the performance on the 3T-VAR task in comparison with both the 2T-VAR and 3T-FIX tasks. A difference between only one of these comparisons would suggest a “cross-track” effect of the variable distractor independent of whether it was in the 2T or 3T condition, or impairment of 3T discriminations independent of whether the distractor was presented at a fixed or variable interval. Thus, we performed a two-way ANOVA to determine if there was an interaction between the 2T/3T and FIX/VAR factors.

Supplemental Data

The Supplemental Data for this article can be found online at <http://www.neuron.org/cgi/content/full/53/3/427/DC1>.

ACKNOWLEDGMENTS

This research was supported by the NIH (MH60163). We would like to thank Jennifer Bramen, Allison Doupe, and Ladan Shams for helpful comments on this manuscript, and the many volunteers who participated in these studies.

Received: April 26, 2006

Revised: September 7, 2006

Accepted: January 2, 2007

Published: January 31, 2007

REFERENCES

- Allan, L.G. (1979). The perception of time. *Percept. Psychophys.* *26*, 340–354.
- Brody, C.D., Hernandez, A., Zainos, A., Lemus, L., and Romo, R. (2002). Analysing neuronal correlates of the comparison of two sequentially presented sensory stimuli. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* *357*, 1843–1850.
- Brody, C.D., Hernandez, A., Zainos, A., and Romo, R. (2003). Timing and neural encoding of somatosensory parametric working memory in macaque prefrontal cortex. *Cereb. Cortex* *13*, 1196–1207.
- Brown, S.W. (1997). Attentional resources in timing: interference effects in concurrent temporal and nontemporal working memory tasks. *Percept. Psychophys.* *59*, 1118–1140.
- Buhusi, C.V., and Meck, W.H. (2005). What makes us tick? Functional and neural mechanisms of interval timing. *Nat. Rev. Neurosci.* *6*, 755–765.
- Buonomano, D.V. (2000). Decoding temporal information: a model based on short-term synaptic plasticity. *J. Neurosci.* *20*, 1129–1141.
- Buonomano, D.V. (2005). A learning rule for the emergence of stable dynamics and timing in recurrent networks. *J. Neurophysiol.* *94*, 2275–2283.
- Buonomano, D.V., and Karmarkar, U.R. (2002). How do we tell time? *Neuroscientist* *8*, 42–51.
- Buonomano, D.V., and Merzenich, M.M. (1995). Temporal information transformed into a spatial code by a neural network with realistic properties. *Science* *267*, 1028–1030.
- Carr, C.E. (1993). Processing of temporal information in the brain. *Annu. Rev. Neurosci.* *16*, 223–243.
- Church, R.M. (1984). Properties of the internal clock. In *Annals of the New York Academy of Sciences*, volume 423, *Timing and Time Perception*, J. Gibbon and L. Allan, eds. (New York: New York Academy of Sciences), pp. 556–582.
- Coull, J.T., Vidal, F., Nazarian, B., and Macar, F. (2004). Functional anatomy of the attentional modulation of time estimation. *Science* *303*, 1506–1508.
- Creelman, C.D. (1962). Human discrimination of auditory duration. *J. Acoust. Soc. Am.* *34*, 582–593.
- Durstewitz, D. (2003). Self-organizing neural integrator predicts interval times through climbing activity. *J. Neurosci.* *23*, 5342–5353.
- Ferster, D., and Miller, K.D. (2000). Neural mechanisms of orientation selectivity in the visual cortex. *Annu. Rev. Neurosci.* *23*, 441–471.
- Gibbon, J. (1977). Scalar expectancy theory and Weber's law in animal timing. *Psychol. Rev.* *84*, 279–325.
- Gibbon, J., Malapani, C., Dale, C.L., and Gallistel, C.R. (1997). Toward a neurobiology of temporal cognition: advances and challenges. *Curr. Opin. Neurobiol.* *7*, 170–184.
- Gilbert, C.D., Sigman, M., and Christ, R.E. (2000). The neural basis of perceptual learning. *Neuron* *31*, 681–697.
- Grondin, S. (2001). From physical time to the first and second moments of psychological time. *Psychol. Bull.* *127*, 22–44.
- Grondin, S., and Rammsayer, T. (2003). Variable foreperiods and temporal discrimination. *Q. J. Exp. Psychol. A.* *56*, 731–765.
- Grondin, S., and Rousseau, R. (1991). Judging the duration of multimodal short empty time intervals. *Percept. Psychophys.* *49*, 245–256.
- Hahnloser, R.H.R., Kozhevnikov, A.A., and Fee, M.S. (2002). An ultra-sparse code underlies the generation of neural sequences in a songbird. *Nature* *419*, 65–70.
- Hicks, R.E., Miller, G.W., and Kinsbourne, M. (1976). Prospective and retrospective judgments of time as a function of amount of information processed. *Am. J. Psychol.* *89*, 719–730.
- Hubel, D.H., and Wiesel, T.N. (1962). Receptive fields, binocular interaction and functional architecture in the cat's visual cortex. *J. Physiol.* *160*, 106–154.
- Ivry, R. (1996). The representation of temporal information in perception and motor control. *Curr. Opin. Neurobiol.* *6*, 851–857.
- Ivry, R.B., and Spencer, R.M.C. (2004). The neural representation of time. *Curr. Opin. Neurobiol.* *14*, 225–232.
- Janssen, P., and Shadlen, M.N. (2005). A representation of the hazard rate of elapsed time in the macaque area LIP. *Nat. Neurosci.* *8*, 234–241.
- Johnston, A., Arnold, D.H., and Nishida, S. (2006). Spatially localized distortions of event time. *Curr. Biol.* *16*, 472–479.
- Karmarkar, U., and Buonomano, D.V. (2003). Temporal specificity of perceptual learning in an auditory discrimination task. *Learn. Mem.* *10*, 141–147.
- Kilgard, M.P., and Merzenich, M.M. (2002). Order-sensitive plasticity in adult primary auditory cortex. *Proc. Natl. Acad. Sci. USA* *99*, 3205–3209.
- Knüsel, P., Wyss, R., König, P., and Verschure, P.F.M.J. (2004). Decoding a temporal population code. *Neural Comput.* *16*, 2079–2100.
- Leon, M.I., and Shadlen, M.N. (2003). Representation of time by neurons in the posterior parietal cortex of the macaque. *Neuron* *38*, 317–327.
- Levitt, H. (1971). Transformed up-down methods in psychoacoustics. *J. Acoust. Soc. Am.* *49*, 467–477.
- Lewis, P.A., and Miall, R.C. (2003). Distinct systems for automatic and cognitively controlled time measurements: evidence from neuroimaging. *Curr. Opin. Neurobiol.* *13*, 250–255.
- Maass, W., Natschläger, T., and Markram, H. (2002). Real-time computing without stable states: a new framework for neural computation based on perturbations. *Neural Comput.* *14*, 2531–2560.
- Macar, F., Grondin, S., and Casini, L. (1994). Controlled attention sharing influences time estimation. *Mem. Cognit.* *22*, 673–686.
- Macar, F., Lejeune, H., Bonnet, M., Ferrara, A., Pouthas, V., Vidal, F., and Maquet, P. (2002). Activation of the supplementary motor area and of attentional networks during temporal processing. *Exp. Brain Res.* *142*, 475–485.
- Markram, H., Wang, Y., and Tsodyks, M. (1998). Differential signaling via the same axon of neocortical pyramidal neurons. *Proc. Natl. Acad. Sci. USA* *95*, 5323–5328.
- Martin, S.J., Grimwood, P.D., and Morris, R.G.M. (2000). Synaptic plasticity and memory: an evaluation of the hypothesis. *Annu. Rev. Neurosci.* *23*, 649–711.
- Matell, M.S., and Meck, W.H. (2004). Cortico-striatal circuits and interval timing: coincidence detection of oscillatory processes. *Brain Res. Cogn. Brain Res.* *21*, 139–170.
- Mauk, M.D., and Buonomano, D.V. (2004). The neural basis of temporal processing. *Annu. Rev. Neurosci.* *27*, 304–340.
- Mauk, M.D., and Ruiz, B.P. (1992). Learning-dependent timing of Pavlovian eyelid responses: differential conditioning using multiple interstimulus intervals. *Behav. Neurosci.* *106*, 666–681.
- Meck, W.H. (1984). Attentional bias between modalities: Effect on the internal clock, memory, and decision stages used in animal time discrimination. In *Annals of the New York Academy of Sciences*, volume 423, *Timing and Time Perception*, J. Gibbon and L. Allan, eds. (New York: New York Academy of Sciences), pp. 528–541.
- Meck, W.H. (1996). Neuropharmacology of timing and time perception. *Brain Res. Cogn. Brain Res.* *3*, 227–242.
- Medina, J.F., and Mauk, M.D. (2000). Computer simulation of cerebellar information processing. *Nat. Neurosci.* *3*, 1205–1211.
- Medina, J.F., Carey, M.R., and Lisberger, S.G. (2005). The representation of time for motor learning. *Neuron* *45*, 157–167.

- Meegan, D.V., Aslin, R.N., and Jacobs, R.A. (2000). Motor timing learned without motor training. *Nat. Neurosci.* *3*, 860–862.
- Miall, C. (1989). The storage of time intervals using oscillating neurons. *Neural Comput.* *1*, 359–371.
- Miller, K.D., Keller, J.B., and Stryker, M.P. (1989). Ocular dominance column development: analysis and simulation. *Science* *245*, 605–615.
- Morrone, M.C., Ross, J., and Burr, D. (2005). Saccadic eye movements cause compression of time as well as space. *Nat. Neurosci.* *8*, 950–954.
- Niki, H., and Watanabe, M. (1979). Prefrontal and cingulate unit activity during timing behavior in the monkey. *Brain Res.* *171*, 213–224.
- Rammsayer, T.H. (1999). Neuropharmacological evidence for different timing mechanisms in humans. *Q. J. Exp. Psychol.* *52B*, 273–286.
- Rammsayer, T.H., and Lima, S.D. (1991). Duration discrimination of filled and empty auditory intervals: cognitive and perceptual factors. *Percept. Psychophys.* *50*, 565–574.
- Rammsayer, T.H., and Ulrich, R. (2001). Counting models of temporal discrimination. *Psychon. Bull. Rev.* *8*, 270–277.
- Rao, S.M., Mayer, A.R., and Harrington, D.L. (2001). The evolution of brain activation during temporal processing. *Nat. Neurosci.* *4*, 317–323.
- Reutimann, J., Yakovlev, V., Fusi, S., and Senn, W. (2004). Climbing neuronal activity as an event based cortical representation of time. *J. Neurosci.* *24*, 3295–3303.
- Reyes, A., and Sakmann, B. (1998). Developmental switch in the short-term modification of unitary. *J. Neurosci.* *19*, 3827–3835.
- Schoups, A., Vogels, R., Qian, N., and Orban, G. (2001). Practicing orientation identification improves orientation coding in V1 neurons. *Nature* *412*, 549–553.
- Shannon, R.V., Zeng, F.G., Kamath, V., Wygonski, J., and Ekelid, M. (1995). Speech recognition with primarily temporal cues. *Science* *270*, 303–304.
- Treisman, M. (1963). Temporal discrimination and the indifference interval: implications for a model of the 'internal clock'. *Psychol. Monogr.* *77*, 1–31.
- Wang, X., Lu, T., Snider, R.K., and Liang, L. (2005). Sustained firing in auditory cortex evoked by preferred stimuli. *Nature* *435*, 341–345.
- Wright, B.A., Buonomano, D.V., Mahncke, H.W., and Merzenich, M.M. (1997). Learning and generalization of auditory temporal-interval discrimination in humans. *J. Neurosci.* *17*, 3956–3963.
- Yang, T., and Maunsell, J.H. (2004). The effect of perceptual learning on neuronal responses in monkey visual area V4. *J. Neurosci.* *24*, 1617–1626.
- Zakay, D., and Tsal, Y. (1989). Awareness of attention allocation and time estimation accuracy. *Bull. Psychon. Soc.* *27*, 209–210.
- Zhang, Z. (2004). Maturation of Layer V pyramidal neurons in the rat prefrontal cortex: intrinsic properties and synaptic function. *J. Neurophysiol.* *91*, 1171–1182.
- Zucker, R.S. (1989). Short-term synaptic plasticity. *Annu. Rev. Neurosci.* *12*, 13–31.